

PROTOCOLS

RABBIT 55.—Subcutaneous injection of 0.015 c. c. of dichlorethylsulphide. Very slight general reaction, with apparently complete recovery, except for the local lesion, 26 days after injection.

Date	Temperature (degrees)	Weight, in grams	Notes
Dec. 16	102.3	1,870	0.015 c. c. dichlorethylsulphide subcutaneously.
17	102.5	1,750	No diarrhoea.
18	102.7	1,740	Do.
19	102.6	1,870	Do.
20	103.1	1,880	Do.
21	102.6	1,930	Do.
22	102.6	1,950	Slight diarrhoea.
23	103.2	2,010	No diarrhoea.
24	103.5	1,970	Do.
25	102.6	2,010	Do.
26	102.6	2,060	Do.
27	103.0	2,100	Do.
28	103.3	2,200	Do.
29	104.1	2,090	Do.
30	103.4	2,200	Do.
31	103.1	2,130	Do.
Jan. 1	102.8	2,120	Do.
2	103.3	2,070	Do.
3	103.0	2,140	Do.
4	102.6	2,150	Do.
5	103.3	2,130	Do.
6	102.8	2,090	Do.
7	103.0	2,090	Do.
8	103.4	2,090	Do.
9	103.5	2,150	Do.
10	102.2	2,300	Do.
11	102.8	2,190	No diarrhoea. Lesion is now nearly healed. There is a hairless area about 1 cm. in diameter, which is covered with a shiny smooth cicatrix except for a small area in its central portion.

RABBIT 48.—Subcutaneous injection of 0.03 c. c. of dichlorethylsulphide. Slight general reaction. Living, 37 days after injection, with apparent complete recovery, except for the local lesion.

Date	Temperature (degrees)	Weight, in grams	Notes
Dec. 5	101.6	2,100	0.03 c. c. dichlorethylsulphide subcutaneously. No signs of local irritation.
6	102.6	2,020	No diarrhoea. Animal very thirsty.
7	102.0	2,050	Do.
8	103.0	2,060	Do.
9	101.0	1,990	Do.
10	101.8	1,940	Slight diarrhoea.
11	103.6	1,950	No evidence of diarrhoea.
12	102.4	1,990	Perianal hair somewhat soiled.
13	103.6	1,900	No diarrhoea.
14	104.4	1,870	Do.
15	103.9	2,000	Do.
16	103.2	2,030	Do.
17	102.7	1,930	Do.
18	102.6	1,830	Do.
19	101.6	1,810	Do.
20	102.6	2,040	Do.
21	103.6	2,000	Do.
22	102.2	2,080	Do.
23	102.1	2,110	Do.
24	102.7	2,090	Do.
25	103.3	2,000	Do.
26	101.6	2,060	Do.
27	102.0	2,070	Do.
28	102.3	2,150	Do.
29	103.5	2,110	Do.
30	103.7	2,140	Do.
31	103.6	2,170	Do.
Jan. 1	103.1	2,020	Do.
2	102.9	2,030	Do.
3	102.8	2,250	Do.
4	103.5	2,170	Do.
5	103.2	2,060	Do.
6	103.4	2,020	Do.
7	103.2	1,990	Do.
8	102.1	2,020	Do.
9	102.6	2,060	Do.
10	102.8	2,240	Do.
11	102.6	2,100	No diarrhoea. The local lesion is not yet completely healed. There is an indurated area about 1 by 1.5 cm. over which there is an incomplete regeneration of epithelium. This area is firmly adherent to the underlying tissue.