

APPENDIX 18

Ratio of Battle Wounds to Combat Exhaustion

First U. S. Army Divisions in France

June and July 1944

DIVISION	JUNE			JULY		
	Wounded	Exhaustion	Ratio	Wounded	Exhaustion	Ratio
1st Infantry	954	161	5.9 to 1	477	94	5.1 to 1
2nd Infantry	1909	361	5.3 to 1	2086	471	4.4 to 1
2nd Armored	127	29	4.4 to 1	1125	283	4.0 to 1
3rd Armored	325	24	13.5 to 1	1150	387	3.0 to 1
4th Armored	—	—	—	299	172	1.7 to 1
4th Infantry	4380	462	9.5 to 1	2212	610	3.6 to 1
5th Infantry	—	—	—	1066	128	8.4 to 1
8th Infantry	—	—	—	2115	282	7.5 to 1
9th Infantry	1900	138	13.8 to 1	3252	471	6.9 to 1
28th Infantry	—	—	—	—	—	—
29th Infantry	2053	287	7.2 to 1	2791	1027	2.7 to 1
30th Infantry	345	40	8.6 to 1	4563	1309	3.5 to 1
35th Infantry	—	—	—	2531	1106	2.3 to 1
79th Infantry	1179	140	8.4 to 1	2706	608	4.5 to 1
82nd Airborne	2223	179	12.4 to 1	725	70	10.4 to 1
83rd Infantry	61	3	20.3 to 1	5295	256	20.7 to 1
90th Infantry	1537	299	5.2 to 1	3114	578	5.4 to 1
101st Airborne	1536	80	19.3 to 1	—	6	—
Total all Divisions	18529	2203	8.4 to 1	35506	7858	4.5 to 1

Source : Combat Medical Statistical Reports ETOUSA MD Forms 323.