

TABLE 60.—AVERAGE PLASMA NITROGENOUS WASTE PRODUCTS AND PHOSPHORUS
IN PATIENTS WHO DIED OF UREMIA

Days after Wounding	NPN	Creatinine	Uric Acid	Phosphorus
	<i>mg. per 100 cc.</i>			
First	69±10.3 (9 cases)	3.0±0.24 (9 cases)	4.1 (3 cases)	4.2±0.47 (8 cases)
Second	118±11.5 (13 cases)	4.8±0.24 (13 cases)	7.3±1.1 (5 cases)	4.1±0.28 (12 cases)
Third	132±6.2 (16 cases)	6.3±0.26 (16 cases)	12.8±2.4 (8 cases)	5.8±0.42 (14 cases)
Fourth	178±5.1 (16 cases)	8.1±0.47 (16 cases)	12.9 (3 cases)	5.9±0.36 (14 cases)
Fifth	245±18.9 (13 cases)	8.9±0.53 (13 cases)	15.0 (3 cases)	7.1±0.96 (12 cases)
Sixth	253±13.0 (8 cases)	9.5±0.70 (9 cases)	18.4 (3 cases)	6.9±0.22 (8 cases)
Seventh	234±26.3 (4 cases)	7.8±1.83 (4 cases)	6.6 (3 cases)
Eighth	255 (2 cases)	11.3 (2 cases)	7.1 (2 cases)
Ninth	278±16.0 (5 cases)	11.4±0.44 (5 cases)	6.8 (3 cases)
Tenth	314 (3 cases)	11.4 (3 cases)	7.7 (3 cases)
Eleventh	296 (2 cases)	10.3 (2 cases)	6.8 (2 cases)

Note: Data are given on this table for days post-wounding rather than days postoperative as in the preceding table.