

**Table 47.—Fractured tibia, United States veterans of the World War, rated 10–29 per cent disabled on first examination, showing interval elapsing between injury and last rating and degree of disability on last rating, as of January 1, 1938 \***

Months (interval elapsing between injury and last rating)	Degree of impairment on last rating †										Total ‡						
	10-19		20-29		30-39		40-49		50-59		70-79		100	Less than 10 per cent	Temporary	Permanent	Total §
	Temporary	Permanent	Temporary	Permanent	Temporary	Permanent	Temporary	Permanent	Temporary	Permanent	Temporary	Permanent					
1																	
2	4		4														1
3			3														3
4																	
5																	
6																	
7																	
8																	
9																	
10																	
11																	
12																	
<b>Total</b>	<b>20</b>	<b>0</b>	<b>14</b>	<b>0</b>									<b>14</b>	<b>36</b>	<b>12</b>		<b>60</b>
1																	
2		2											1	0	0		1
3		1											1	0	0		1
4		2											1	0	0		1
5		3											1	0	0		1
6		5											2	0	0		2
7		5											3	0	0		3
8		5											3	0	0		3
9		3											2	0	0		2
10		3											2	0	0		2
11		3											2	0	0		2
12		3											2	0	0		2
<b>Total</b>	<b>31</b>	<b>35</b>	<b>59</b>	<b>12</b>	<b>2</b>	<b>0</b>			<b>1</b>			<b>2</b>	<b>126</b>	<b>123</b>	<b>48</b>		<b>300</b>
1																	
2		20											1	0	0		1
3		24											2	0	0		2
4		14											1	0	0		1
5		10											1	0	0		1
6		10											1	0	0		1
7		14											2	0	0		2
8		17											3	0	0		3
9		17											3	0	0		3
10		17											3	0	0		3
11		17											3	0	0		3
12		17											3	0	0		3
<b>Total</b>	<b>67</b>	<b>636</b>	<b>35</b>	<b>125</b>	<b>3</b>	<b>14</b>	<b>1</b>	<b>10</b>	<b>2</b>	<b>4</b>	<b>1</b>	<b>4</b>	<b>188</b>	<b>113</b>	<b>591</b>		<b>892</b>
1																	
2		26											1	0	0		1
3		20											2	0	0		2
4		14											1	0	0		1
5		13											1	0	0		1
6		17											2	0	0		2
7		17											2	0	0		2
8		17											2	0	0		2
9		17											2	0	0		2
10		17											2	0	0		2
11		17											2	0	0		2
12		17											2	0	0		2
<b>Total</b>	<b>159</b>	<b>80</b>	<b>12</b>	<b>35</b>	<b>1</b>	<b>12</b>		<b>9</b>	<b>4</b>	<b>1</b>	<b>1</b>	<b>2</b>	<b>10</b>	<b>36</b>	<b>47</b>	<b>136</b>	<b>219</b>
1																	
2																	
3																	
4																	
5																	
6																	
7																	
8																	
9																	
10																	
11																	
12																	
<b>Total</b>	<b>1</b>	<b>1</b>											<b>1</b>				<b>1</b>
<b>Grand total</b>	<b>187</b>	<b>596</b>	<b>69</b>	<b>161</b>	<b>6</b>	<b>27</b>	<b>1</b>	<b>19</b>	<b>6</b>	<b>6</b>	<b>3</b>	<b>2</b>	<b>16</b>	<b>209</b>	<b>137</b>	<b>768</b>	<b>1,474</b>

\* Source of information: Coordination Service, Evaluation Division, U. S. Veterans' Bureau.

† Where the column is omitted there were no cases.

‡ Exclusive of patients who died during the six-year period of the study.