

TABLE 52.—Fractured tibia and fibula among United States veterans of the World War, rated less than 10 per cent disabled on first examination, showing interval elapsing between injury and last rating and degree of disability on last rating, January 1, 1926 \*

Months (interval elapsing between injury and last rating)	Degree of impairment on last rating †										Total									
	10-19		20-29		30-39		40-49		50-59		70-79		80-89		100		Less than 10 per cent	Temporary	Permanent	Total ‡
	Temporary	Permanent	Temporary	Permanent	Temporary	Permanent	Temporary	Permanent	Temporary	Permanent	Temporary	Permanent	Temporary	Permanent						
3																	25			25
6																	4			4
9																	1			1
12																	3			3
15																	3			3
18																	4			4
21																	4			4
24																	4			4
Total																	52			52
27																	4		1	5
30																	16			16
33																	26			26
36																	17			17
39		1	1														17	1	1	19
42																	1			1
45		1		1													16	2	1	19
48																	12			12
49																	1		2	3
Total		3	1	1			1	2									109	4	6	119
51																	9			9
54		1	3														8	2	3	13
57		1	3														7	1	4	12
60																	2	2	2	6
63																	7	1	1	9
66		1	1														11	2	2	15
69																	9	1	4	14
72																	2	2	2	6
75																	8	1	3	12
78																	1			1
81																	2			2
84																	3			3
87																	3			3
90																	1		3	4
93																	1			1
Total		1	12		1	3	2	4									12	4	21	36
Grand total		9	36		3	5	4	10		1	1	1	1	1	3	3	235	19	34	309

\* Source of information: Coordination Service, Evaluation Division, U. S. Veterans' Bureau.

† Where the column is omitted there were no cases.

‡ Exclusive of patients who died during the six-year period of the study.