

TABLE 54.—Fractured tibia and fibula, United States veterans of the World War, rated 30-49 per cent disabled on first examination, showing interval elapsing between injury and last rating and degree of disability on last rating, as of January 1, 1926 *

Months (interval elapsing between injury and last rating)	Degree of impairment on last rating *												Total						
	10-19		20-29		30-39		40-49		50-59		60-69	70-79	100	* Less than 10 per cent	Temporary	Permanent	Total *		
	Temporary	Permanent	Temporary	Permanent	Temporary	Permanent	Temporary	Permanent	Temporary	Permanent	Temporary	Permanent							
12.....							1									1	1		
15.....				1	2		1	1								3	3		
21.....							1	2								3	3		
24.....	1															1	1		
Total.....	1			1	2		3	3								1	6	4	
27.....						3		1	1								4	1	
30.....		1		2				1	1							2	4	1	
33.....	1			1		2										2	3	2	
36.....				3	3			2	2							2	4	2	
39.....	1			1	1							1				1	2	1	
42.....	1			3	1	2				1						1	3	2	
45.....	1			1	2	1										1	1	1	
48.....		2		4	1					1						2	3	1	
Total.....	6	3	4	13	11	10	1	25		2			1			9	25	53	45
51.....	1	1	2	1	1	1	5	1	8				1			2	6	15	23
54.....		6		5	1		1	1	5							1	2	17	20
57.....				6			6	1	11							2	1	25	28
60.....	2			1	6		8	6	10				3			1	6	30	35
63.....	1	13	2	9	3		6		17				2			2	6	47	55
66.....		8	1	13	1	14	7	7	7		3	1				1	4	46	47
69.....		3	3	5		8	1	17	10				1			3	3	42	45
72.....						8		10			1		3			6	27	33	33
Total.....	4	45	9	54	4	55	4	85		6	3		10			8	31	149	206
75.....		8		2	1	2				4	1					1	2	26	29
78.....	1			8	1		1	3								3	15	16	16
81.....		2		3		2		4	1	2						1	3	13	17
84.....		1	1	1				3		1			2			4	6	10	10
87.....	1			1				3	1	2			1			3	3	6	6
90.....				1												1	1	1	1
93.....								1								1	1	1	1
Total.....	2	12	1	16	2	6	1	22	2	9	2	2	4			1	14	67	82
96.....								2										2	2
Grand total.....	13	61	14	84	19	71	9	137	2	17	5	2	15			19	74	373	406

* Source of information: Coordination Service, Evaluation Division, U. S. Veterans' Bureau.

* Where the column is omitted there were no cases.

* Exclusive of patients who died during the six-year period of the study.