

TABLE 74.—Fractures of the long bones, United States veterans of the World War, showing the number and percentage of cases which reached their stationary level after periods of 2, 3, 4, 5, or more years, as of January 1, 1926

Bone	Total		Unclassified		Period									
					2 years		3 years		4 years		5 years		Over 5 years	
	Cases	Per cent	Cases	Per cent	Cases	Per cent	Cases	Per cent	Cases	Per cent	Cases	Per cent	Cases	Per cent
Femur.....	5, 138	100. 00			259	5. 04	455	8. 47	637	12. 40	1, 185	23. 06	2, 622	51. 03
Tibia.....	2, 299	100. 00	85	3. 74	118	5. 16	212	9. 26	262	11. 42	478	20. 88	1, 129	49. 32
Fibula.....	1, 243	100. 00	203	16. 33	89	7. 16	123	9. 90	158	12. 31	221	17. 78	454	36. 52
Tibia and fibula.....	4, 482	100. 00	23	. 51	226	5. 04	433	10. 10	562	12. 53	1, 027	22. 90	2, 194	48. 92
Humerus.....	4, 328	100. 00			214	4. 94	303	7. 00	462	11. 37	934	21. 56	2, 365	54. 64
Ulna.....	1, 369	100. 00	169	14. 49	60	5. 13	83	7. 10	129	11. 64	230	19. 67	498	42. 60
Radius.....	1, 356	100. 00	101	7. 45	110	8. 11	109	8. 04	143	10. 69	306	22. 53	591	43. 58
Radius and ulna.....	2, 340	100. 00	20	. 85	132	5. 64	185	7. 91	311	13. 29	538	22. 99	1, 154	49. 32
Total.....	22, 348	100. 00	602	2. 69	1, 208	5. 41	1, 903	8. 52	2, 695	12. 06	4, 933	22. 07	11, 007	49. 25