

TABLE 16.—BLOOD CHANGES IN 69 SEVERELY WOUNDED MEN CLASSIFIED
ACCORDING TO DEGREE OF SHOCK

A. No Shock—12 Patients

Case No.	Blood Volume and Total Hemoglobin Determinations <i>Percent deviation from normal</i>						Blood Pressure at Time of Blood Volume Determina- tion <i>mm. Hg</i>	Time after Wound- ing <i>hours</i>	Blood Sugar <i>mg. per 100 cc.</i>
	Blood Volume			Hemoglobin					
	From Ob- served Value	By Correction		From Ob- served Value	By Correction				
		A	B		A	B			
91	-26.0	-37.7	-37.7	-43.1	-52.9	-52.9	108/78	2¾	—
102	-21.3	-36.9	-29.1	-31.1	-39.0	-39.0	150/80	3¼	—
*124	+15.3	-2.8	-2.8	+14.7	-1.6	-1.6	120/64	10¼	—
134	-7.8	-11.3	-7.8	-3.8	-3.8	-3.8	120/76	15¼	—
A-4	-8.2	-8.2	-8.2	-31.1	-31.1	-31.1	132/72	4¼	93
A-13	-18.4	-18.4	-18.4	-36.2	-36.2	-36.2	150/70	18¾	121
A-16	-8.2	-10.5	-8.2	-12.2	-12.2	-12.2	130/70	5¼	139
A-19	-31.9	-31.9	-31.9	-42.5	-42.5	-42.5	116/78	8¼	167
A-22	-2.8	-2.8	-2.8	+11.5	+11.5	+11.5	140/80	1½	110
A-23	+17.0	+17.0	+17.0	+13.3	+13.3	+13.3	128/80	4½	127
A-24	-0.3	-10.3	-10.3	+4.1	-6.4	-6.4	110/70	4½	140
A-39	-7.1	-7.1	-7.1	-20.5	-20.5	-20.5	124/72	4	184

* Case of crush injury.